






























































	CLASS/DRUG	USUAL ADULT DAILY DOSING [†]	SPECIAL CONSIDERATIONS	ACTUAL SIZE
NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (NNRTI)	VIRAMUNE® (nevirapine)	1 x 200 mg tablet 2 times a day 	Starting Dose: 1 x 200 mg tablet once daily for first 14 days.	
	Stocrin® (efavirenz)	1 x 600 mg tablet once daily 	Take at bedtime. Take away from meals.	
	Intelence® (etravirine)	2 x 100 mg tablets 2 times a day 	Take after a meal.	
NUCLEOSIDE / NUCLEOTIDE REVERSE TRANSCRIPTASE INHIBITORS (NRTI / NtRTI)	Zerit® (stavudine-also known as d4T)	1 x 40 mg capsule 2 times a day 1 x 30 mg capsule 2 times a day  or 		
	Retrovir® (zidovudine-also known as AZT)	2 x 100 mg capsules 3 times a day 1 x 250 mg capsule 2 times a day  or 		
	3TC® (lamivudine)	1 x 150 mg tablet 2 times a day 1 x 300 mg tablet once daily  or 		
	Combivir® (lamivudine/zidovudine)	1 x 150mg / 300 mg tablet 2 times a day 		
	Ziagen® (abacavir)	1 x 300 mg tablet 2 times a day 2 x 300 mg tablets once daily  or 		
	Kivexa® (abacavir/lamivudine)	1 x 600 mg / 300 mg tablet once daily 		
	Trizivir® (abacavir/lamivudine/zidovudine)	1 x 300 mg / 150 mg / 300 mg tablet 2 times a day 		
	Videx® EC (didanosine-also known as ddl)	1 x 400 mg capsule once daily 1 x 250 mg capsule once daily  or 	Take on an empty stomach.	
	Emtriva® (emtricitabine-also known as FTC)	1 x 200 mg capsule once daily 		
	Viread® (tenofovir disoproxil fumarate)	1 x 300 mg tablet once daily 	Preferably taken with a meal.	
	Truvada® (tenofovir disoproxil fumarate/emtricitabine)	1 x 300 mg / 200 mg once daily 	Preferably taken with a meal.	
PROTEASE INHIBITORS (PI)	Crixivan® (indinavir)	2 x 400 mg capsules 3 times a day 	Take on an empty stomach. Drink at least 1.5 L of liquid daily. Lower daily doses may be used with other PIs.	
	Invirase® (saquinavir)	2 x 500 mg tablets 2 times a day 	Take each 2 x 500 mg dose with 100mg ritonavir. Take within 2 hours of a full meal.	
	Kaletra® (lopinavir/ritonavir)	2 x 200mg/50mg tablets 2 times a day 4 x 200mg/50mg tablets once daily  or 	Can be taken with or without food.	
	Telzir® (fosamprenavir calcium)	1 x 700 mg tablet 2 times a day 2 x 700 mg tablets once daily  or 	Take each tablet with 100 mg ritonavir. Can be taken with or without food.	
	Reyataz® (atazanavir)	1 x 300 mg capsule once daily 2 x 200 mg capsules once daily  or 	Take each 1 x 300 mg dose with 100 mg ritonavir. Take with food.	
	APTIVUS® (tipranavir)	2 x 250 mg capsules 2 times a day 	Take each 2 x 250 mg dose with 200 mg ritonavir. Take with food.	
	Prezista® (darunavir)	2 x 300 mg tablet 2 times a day 	Take each 2 x 300 mg dose with 100 mg ritonavir. Take with food.	
	Norvir® (ritonavir)	Mainly used in small doses (100 mg / 200 mg) in combination with other PIs	Take with food.	
ENTRY INHIBITORS	Fuzeon® (enfuvirtide-also known as T-20)	1 x 90 mg vial 2 times a day 	Give by subcutaneous injection.	 (Not actual size)
	Celsentri® (maraviroc)	1 x 150 mg tab or 1 x 300 mg tab or 2 x 300 mg tabs 2 times a day 2 times a day 2 times a day   	Dose is dependent on concomitant medication.	
INTEGRASE INHIBITORS	Isentress® (raltegravir)	1 x 400 mg tablet 2 times a day 		
COMBINED AGENTS	Atripla® (tenofovir disoproxil fumarate/emtricitabine/efavirenz)	1 x 300 mg / 200 mg / 600 mg tablet once daily 	Take on an empty stomach. Take at bedtime.	

TM = Trademark © Registered Trademark † Please refer to product information before prescribing as dosing may change with some combinations or patient types *Or as directed by physician